

## Inhibition Kinetic Spectrophotometric Determination of Resorcinol Based on Its Inhibitory Effect on Formaldehyde Catalyzed Reaction Between Bromate and Spadns

MOHSEN KEYVANFARD

Faculty of Science, Islamic Azad University-Majlesi Branch, Isfahan, Iran  
E-mail: keyvan45638@yahoo.com

A new, sensitive, simple, inexpensive and fast kinetic spectrophotometric method was developed for the determination of trace amounts of resorcinol over the range of 0.02-1.40  $\mu\text{g/mL}$ . The method is based on the inhibitory effect of resorcinol on the formaldehyde catalyzed oxidation reaction of of spadns by bromate in acidic media is reported. The reaction was monitored spectrophotometrically by measuring the decrease in absorbance of spadns at 530 nm with a fixed-time 0.5-3.0 min from initiation of the reaction. The detection limit is 15.2 ng/mL and relative standard deviation of 0.4 and 1.2  $\mu\text{g/mL}$  resorcinol for 5 replicate measurements was 2.2 and 2.7 %, respectively. The method was applied to the determination of resorcinol in water samples.

**Key Words:** Resorcinol, Kinetic, Determination, Spadns, Inhibitory.

### INTRODUCTION

Resorcinol is an important industrial chemical material used widely in the fields of rubber, plastics and organic synthesis industries, wood adhesives, fire retardants and UV stabilizer, *etc.* It is derived from various resins and tannins but most commonly by fusing sodium hydroxide with *meta*-benzene-disulfuric acid. Global output of resorcinol in 2004 was about 41.83 million tons. This compound is a moderate toxic substance and easily soluble in water<sup>1</sup>. There is a growing need for developing highly sensitive, simple methods to detect resorcinol in the wastewater at a low level. The major methods for the determination of resorcinol that have already been reported are high-performance liquid chromatography<sup>2-6</sup> and gas chromatography<sup>7,8</sup>. The separations of these methods are efficient, but require expensive instrument and therefore are expensive. Another resorcinol measurement method is ultraviolet-visible spectrophotometry<sup>9</sup>. This method is convenient but its sensitivity is low. Some of proposed kinetic spectrophotometric method for determination of resorcinol were expensive or their sensitivity are low<sup>10,11</sup>. Therefore, the need for a sensitive, simple, rapid and sensitive kinetic spectrophotometric method for the determination of resorcinol is clearly recognized.

In this paper, we developed and validated a rapid, sensitive kinetic spectrophotometric method for the determination of resorcinol. Here, we report a kinetic method for trace determination of resorcinol, based on its inhibitory effect on the formaldehyde catalyzed oxidation reaction of of spadns by bromate in acidic media.

## EXPERIMENTAL

Doubly distilled water and analytical reagent grade chemicals were used during all of the experimental studies. All glassware were cleaned with detergent solution, rinsed with tap water, soaked in dilute  $\text{HNO}_3$  solution (2 % v/v), rinsed with water and dried. Spadns solution  $7.36 \times 10^{-4}$  M was prepared by dissolving 0.0419 g of the compound (Merck) in water and solution was diluted to the mark in a 100 mL volumetric flask. Bromate stock solution 0.015 M was prepared by dissolving 0.626 g of potassium bromate ( $M = 167$ ) in water and diluting to 250 mL in a 250 mL volumetric flask. An aqueous formaldehyde stock solution,  $1000 \mu\text{g mL}^{-1}$ , was prepared by diluting 2.5 mL of 37 % w/v stock formaldehyde solution to 1 L with water. Standard stock resorcinol solution ( $1000 \mu\text{g/mL}$ ) was prepared by dissolving 0.1 g of resorcinol in water and diluted to 100 mL in a 100-mL volumetric flask. Sulfuric acid solution was prepared by appropriate dilution of concentrated sulfuric acid (Merck). Stock solution ( $1000 \mu\text{g/mL}$ ) of interfering ions were prepared by dissolving suitable salts in water, hydrochloric acid or sodium hydroxide solution.


Absorption spectra were recorded with a CECIL model 7500 spectrophotometer with a 1.0 cm quartz cell. A model 2501 CECIL Spectrophotometer with 1.0 cm glass cuvettes was used to measure the absorbance at a fixed wavelength of at 530 nm. A thermostat water bath (Gallen Kamp Griffin, BGL 240 V) was used to keep the reaction temperature at 27 °C. A stopwatch was used for recording the reaction times.

**Recommended procedure:** All the solutions and distilled water were kept in a thermostated water bath at 27 °C for 20 min for equilibration before starting the experiment. An aliquot of the solution containing 0.20-14.0  $\mu\text{g/mL}$  resorcinol was transferred into a 10 mL volumetric flask and then 1 mL  $7.36 \times 10^{-4}$  M spadns, 2 mL 5 M sulphuric acid and 0.2 mL  $1000 \mu\text{g/mL}$  formaldehyde were added to the flask. The solution was diluted to *ca.* 7 mL with water. Then, 2 mL of 0.015 M bromate was added and the solution was diluted to the mark with water. The solution was mixed and a portion of the solution was transferred to the spectrophotometer cell. The reaction was followed by measuring the decrease in absorbance of the solution against water at 530 nm for 0.5-3.0 min from initiation of the reaction. This signal (sample signal) was labeled as  $\Delta A_s$ . The same procedure was repeated without addition of resorcinol solution and the signal (blank signal) was labeled as  $\Delta A_b$ . Time was measured just after the addition of last drop of bromate solution. Analytical signal was difference between blank signal and sample signal ( $\Delta A_b - \Delta A_s$ ).

## RESULTS AND DISCUSSION

Spadns (I) is a dye that can be oxidized with strong oxidizing agents at slow reaction. Formaldehyde can increasing rate this reaction at ultra-trace level. It is found that trace amount of resorcinol have a inhibitory effect on the this reaction. Therefore, by measuring the decrease in absorbance of spadns for a fixed time of 0.5-3.0 min initiation of the reaction, the resorcinol contents in the sample can be

measured. There are many methods, such as fixed-time, initial rate, rate constant and variable time methods for measuring the kinetic species. Among these, the fixed-time method is the most conventional and simplest, involving the measurement of  $\Delta A$  at 530 nm (Fig. 1).



Structure of spadns (I)




Fig. 1. Absorption spectrum for the resorcinol-spadns- $\text{BrO}_3^-$  system with time. Conditions:  $\text{H}_2\text{SO}_4$ , 1.0 M; spadns,  $7.36 \times 10^{-4}$  M;  $\text{BrO}_3^-$   $3.0 \times 10^{-3}$  M; formaldehyde 20  $\mu\text{g/mL}$ , temperature, 27  $^\circ\text{C}$ ; interval time for each scan, 0.5, 1.5 and 2.5 from initiation of the reaction (a) in presence of 0.4  $\mu\text{g/mL}$  of resorcinol (b) in absence of resorcinol

**Influence of variables:** In order to take full advantage of the procedure, the reagent concentrations must be optimized. The effect of acid concentration, spadns, formaldehyde and bromate concentration and temperature on analytical signal was studied.

The effect of sulfuric acid concentration on the analytical signal was studied in the range of 0.6-1.4 M (Fig. 2). The results show that the analytical signal increases with increasing sulfuric acid concentration up to 1.0 M and decreases at higher concentrations. Therefore, a sulfuric acid concentration of 1.0 M was selected for further study.




Fig. 2. Effect of  $\text{H}_2\text{SO}_4$  concentration on the analytical signal. Conditions spadns,  $5.89 \times 10^{-5}$  M;  $\text{BrO}_3^-$   $3.0 \times 10^{-3}$  M; formaldehyde 20  $\mu\text{g}/\text{mL}$ , temperature, 27  $^\circ\text{C}$  and time of 2.5 min from initiation of the reaction

The influence of spadns concentration on the analytical signal was studied in the concentration range of  $4.41 \times 10^{-5}$  –  $8.83 \times 10^{-5}$  M (Fig. 3). The results show that analytical signal increases with increasing spadns concentration up to  $7.36 \times 10^{-5}$  M and decreases at higher concentrations. Therefore, a spadns concentration of  $7.36 \times 10^{-5}$  M was selected for further study.




Fig. 3. Effect of spadns concentration on the analytical signal. Conditions:  $\text{H}_2\text{SO}_4$ , 1.0 M;  $\text{BrO}_3^-$   $3.0 \times 10^{-3}$  M, formaldehyde 20  $\mu\text{g}/\text{mL}$ , temperature, 27  $^\circ\text{C}$  and time of 2.5 min from initiation of the reaction

Fig. 4 shows the effect of the bromate concentration on the analytical signal for the range of  $1.8 \times 10^{-3} - 3.6 \times 10^{-3}$  M. This analytical signal increases with increasing bromate concentration up to  $3 \times 10^{-3}$  M and decreases at higher concentrations. Therefore, a final concentration of  $3 \times 10^{-3}$  M of bromate was selected as the optimum concentration.




Fig. 4. Influence of  $\text{BrO}_3^-$  concentration on the analytical signal. Conditions:  $\text{H}_2\text{SO}_4$  1.0 M; spadns  $7.36 \times 10^{-4}$  M, formaldehyde 20  $\mu\text{g/mL}$ ; temperature 27 °C and time of 2.5 min from initiation of the reaction

Fig. 5 shows the effect of the formaldehyde concentration on the analytical signal for the range of 10-60  $\mu\text{g mL}^{-1}$ . Analytical signal increases with increasing formaldehyde concentration up to 20  $\mu\text{g mL}^{-1}$  and decreases at higher concentrations. Therefore, a final concentration of 20  $\mu\text{g mL}^{-1}$  of formaldehyde was selected as the optimum concentration.




Fig. 5. Influence of formaldehyde on the analytical signal. Conditions:  $\text{H}_2\text{SO}_4$ , 1 M; spadns  $7.36 \times 10^{-5}$  M;  $\text{BrO}_3^-$   $3.0 \times 10^{-4}$  M; temperature, 27 °C and time of 2.5 min from initiation of the reaction

The effect of the temperature on the analytical signal was studied in the range 20-42 °C with the optimum of the reagents concentrations. The results showed that, as the temperature increases up to 27 °C, the analytical signal increases, whereas higher temperature values decrease the analytical signal ( $\Delta A = \Delta A_b - \Delta A_s$ ). Therefore, 27 °C was selected for further study.

**Calibration graph precision and limit of detection:** Calibration graph were obtained using the fixed-time method. This method was applied to the change in absorbance over an interval of 0.5-3.0 min from initiation of the reaction because it provided the best regression and sensitivity. Under the optimum conditions described above, a linear calibration range 0.02-1.4 µg/mL of resorcinol. The equation of the calibration graph is  $\Delta A = 0.345C + 0.1075$  ( $n = 8$ ,  $r = 0.9998$ ). The calibration graph was constructed by plotting of  $\Delta A = \Delta A_b - \Delta A_s$  at a fixed-time method *versus* resorcinol concentration. The experimental  $3\delta$  limit of detection was 15.2 ng/mL.

The relative standard deviation for five replicate determinations of 0.4 and 1.2 µg/mL resorcinol was 2.2 and 2.7 %, respectively.

**Interference study:** In order to assess the application of the proposed method to synthetic samples, the effect of various ions on the determination of 0.1 µg/mL resorcinol was studied. The tolerance limit was defined as the concentration of added ions causing a relative error less than 3 % (Table-1). The results show that the method is relatively selective for resorcinol determination.

TABLE-1  
EFFECT OF FOREIGN IONS ON THE DETERMINATION OF 0.1 µg/mL RESORCINOL

Species	Tolerance limit ( $w_{ion}/w_{Resorcinol}$ )
Te <sup>4+</sup> , Se <sup>4+</sup> , Na <sup>+</sup> , Mg <sup>2+</sup> , Pb <sup>2+</sup> , C <sub>2</sub> O <sub>4</sub> <sup>2-</sup> , HSO <sub>4</sub> <sup>-</sup> , CO <sub>3</sub> <sup>2-</sup> , HCO <sub>3</sub> <sup>-</sup> , tatarate, borate	1000
Co <sup>2+</sup> , ethanolamine, SO <sub>3</sub> <sup>2-</sup> , S <sub>2</sub> O <sub>8</sub> <sup>2-</sup> , Hg <sup>2+</sup> , Mn <sup>2+</sup> , Fe <sup>2+</sup> , Fe <sup>3+</sup>	500
Ethanol, methanol	400
I <sup>-</sup>	200
Ag <sup>+</sup>	100
SCN <sup>-</sup> , Br <sup>-</sup>	50
S <sub>2</sub> O <sub>3</sub> <sup>2-</sup>	25
Cl <sup>-</sup>	10

**Sample analysis:** In order to evaluate the applicability of the proposed method, water samples were analyzed to determine resorcinol contents. The results are presented in Table-2. Good recoveries with precise results show good reproducibility and accuracy of the method.

TABLE-2  
DETERMINATION OF RESORCINOL IN SYNTHETIC SAMPLES

Sample	Resorcinol added (ng/ml)	Resorcinol found (ng/ml)	Recovery (%)	RSD (n = 5)
Well water	-	-	-	-
Well water	200	214	107	2.2
Well water	800	787	98.3	2.4

## Conclusion

The kinetic-spectrophotometric method developed for the determination of resorcinol is inexpensive, uses readily available reagents, allows rapid determination at low operating costs and shows simplicity, good precision and accuracy compared to other kinetic procedures.

## ACKNOWLEDGEMENT

The author is thankful to the Islamic Azad University-Majlesi Branch for the support of this work.

## REFERENCES

1. A.B. Sangli and V.V. Kanabur, *Environ. Ecol.*, **16**, 642 (1998).
2. G.P. McMahon and M.T. Kelly, *Anal. Chem.*, **70**, 409 (1998).
3. M. Fiehn, *Jekel*, **769**, 189 (1997).
4. H. Kim, H.P. Lee, S. Chung, S. Choi, K. Lee and S. Han, *J. Chromatogr. B*, **792**, 307 (2003).
5. E. Mikami, T. Goto, T. Ohno, H. Matsumoto and M. Nishida, *J. Pharm. Biomed. Anal.*, **28**, 261 (2002).
6. C. Hua, Z. Jian, X. Feng, L. Chun-Ze and W. Guo-Hui, *Anal. Chim. Acta*, **511**, 273 (2004).
7. J. Engeberth, E.A. Schmelz, H.T. Alborn, Y.J. Cardoza, J. Huang and J.H. Tumlinson, *Anal. Biochem.*, **312**, 242 (2003).
8. C. Lartigue-Mattei, C. Lauro-Marty, M. Bastide, J.A. Berger, J.L. Chabard, E. Goutay and J.M. Aiache, *J. Chromatogr. B*, **617**, 140 (1993).
9. B.W. Glombitza and P.C. Schmidt, *J. Pharm. Sci.*, **83**, 751 (1994).
10. H. Parham, N. Pourreza and S. Cheraghi, *Anal. Lett.*, **32**, 1917 (1999).
11. F. Jing, Z. Tao, S. Jianhui and F. Maohong, *J. Fluoresc.*, **17**, 113 (2007).
12. S. Abbasi, M. Esfandyarpour, M.A. Taher and A. Daneshfar, *Spectrochim. Acta A: Mol. Biomol. Spectrosc.*, **67**, 578 (2007).

(Received: 29 January 2010;

Accepted: 22 June 2010)

AJC-8819