

## Synthesis, Characterization and Antibacterial Studies of Transition Metal Complexes of Mannich Base

K. BALAKRISHNAN

Department of Chemistry, Payyanur College, Payyanur,  
P.O. Edat, Distt. Kannur-670 327, India  
E-mail: kbnmanakkat@rediffmail.com


A new series of complexes of Fe(III), Co(II), Ni(II), Cu(I), Zn(II), Hg(II), Pb(II) and Ag(I) with 3-morpholinomethyl-5-( $\beta$ -naphthoxy)-methyl-1,3,4-oxadiazole-2-thione have been synthesized. Characterization of these complexes was carried out by elemental analyses, magnetic susceptibility measurements, thermogravimetric, UV-Vis, IR and  $^1\text{H}$  NMR spectral studies. Though the ligand is potentially polydentate, in the complexes studied, it behaves as a monodentate ligand—bonding either through N(4) of the oxadiazole moiety or through S of the thione group—or as a bidentate bridging ligand. Tetrahedral structures for Fe(III), Co(II), Ni(II), Cu(I), Zn(II), Hg(II) and Pb(II) complexes, tetrahedral polymeric structure for Cd(II) and linear structure for Ag(I) complex have been proposed. Antibacterial studies of the ligand and Fe(III), Co(II), Cu(I) and Zn(II) complexes were also carried out.

**Key Words:** Antibacterial Studies, Mannich base, Transition Metal complexes, Oxadiazoles.

### INTRODUCTION

Substituted 1,3,4-oxadiazole-2-thione and their metal complexes have gained attention due to their wide range of biological activities<sup>1–5</sup>. Some of the 1,3,4-oxadiazole derivatives are found to be acting as corrosion inhibitors in mild steel in acid medium<sup>6</sup>. In the literature, there is only one report on the metal complexes of oxadiazole Mannich bases and their fungicidal activity<sup>7</sup>. Therefore, it was thought worthwhile to synthesize metal complexes of some substituted oxadiazole thione Mannich bases and to investigate their bonding characteristics.

The present communication deals with the synthesis, characterization and biological studies of some metal complexes of 3-morpholinomethyl-5-( $\beta$ -naphthoxy)methyl-1,3,4-oxadiazole-2-thione (L).



(L)

## EXPERIMENTAL

All the chemicals used were of AnalaR or chemically pure grade. Metal salts (chlorides, nitrates and sulphates) and the reagents used were of ExcelaR or SQ quality of Qualigens Fine Chemicals, India. The solvents used were purified by standard methods. Spectroscopic grade solvents were used for spectral studies. The parent oxadiazole, 5-( $\beta$ -naphthoxy)methyl-1,3,4-oxadiazole-2-thione was prepared by the literature method<sup>8,9</sup>. The Mannich base was then prepared by stirring a solution of the thione (0.01 mol), formaldehyde (0.0015 mol) and morpholine (0.01 mol) in 40 mL ethanol for 1 h and keeping overnight. The solid compound formed was separated and washed several times with ethanol. The product was recrystallized from ethanol and then dried. (m.p. = 80°C; yield 70%).

The Fe(III), Co(II), Ni(II), Cu(I), Cd(II) and Hg(II) complexes were prepared by adding a hot ethanolic solution of respective metal chlorides to a refluxing solution of the ligand in ethanol, maintaining the stoichiometric metal-ligand ratios. The solid complexes separated out almost immediately, except in Fe(III) and Ni(II), were filtered off, washed repeatedly with hot ethanol and dried. In the case of Fe(III) and Ni(II), complexes separated out only after raising the pH of the medium by adding sodium acetate (*ca.* 1.50 g) and maintaining the reaction mixture at the refluxing temperature for 0.5 h. For preparing the Zn(II) complex, a hot aqueous solution of Zn(II) sulphate was used, while for Pb(II) and Ag(I) complexes, the starting materials were aqueous solution of respective metal nitrates.

C, H and N contents of the ligand and the complexes were determined microanalytically. Metal, sulphur and chlorine contents were determined by standard procedures<sup>10</sup>. The molar conductances were measured using a Systronics microprocessor-based conductivity meter (model 306, Systronics, India). Magnetic susceptibilities were determined with a Gouy assembly at room temperature, using Hg[Co(NCS)<sub>4</sub>] as the calibrant. Electronic spectra were recorded on a Perkin-Elmer Lambda 15 UV/Vis spectrophotometer in DMF solution. IR spectra were recorded (KBr) on a Perkin-Elmer FT1000 spectrophotometer and <sup>1</sup>H NMR (DMSO-d<sub>6</sub> + CDCl<sub>3</sub>) on a Perkin-Elmer R-32 spectrometer at 90 MHz. Thermograms were recorded on a Perkin-Elmer 7 Series thermal analysis system. The complexes were heated in air at the rate of 10°C per min.

## RESULTS AND DISCUSSION

The ligand was characterized by mass, NMR and IR spectral data. The mass spectrum of the ligand showed molecular ion peak at *m/z*, 357 consistent with the molecular formula C<sub>18</sub>H<sub>19</sub>N<sub>3</sub>O<sub>3</sub>S. The <sup>1</sup>H NMR of the ligand showed signals at  $\delta$  2.76–2.81 (t, 4H —CH<sub>2</sub>—N—CH<sub>2</sub>—),  $\delta$  3.65–3.70 (t, 4H, —CH<sub>2</sub>—O—CH<sub>2</sub>—),  $\delta$  4.99 (s, 2H, —N—CH<sub>2</sub>—N—),  $\delta$  5.16 (s, 2H, —O—CH<sub>2</sub>—) and  $\delta$  7.18–7.82 (m, 7H, Ar—H). IR spectrum of the ligand shows a strong band at 1617 cm<sup>-1</sup> and has been assigned to  $\nu$ (C=N) of oxadiazole ring.  $\nu$ (C=S) of the ligand was assigned at 1122 cm<sup>-1</sup>. In the ligand  $\nu$ (C—O—C) of oxadiazole ring<sup>11</sup> appears at 1432 cm<sup>-1</sup>.

The analytical data support the proposed formulations for the complexes (Table-1). The non-electrolytic nature of the complexes, except the Fe(III) complex, is ascertained by low conductivity values which are much lower than those reported for 1 : 1 electrolytes in the same solvent. For the Fe(III) complex, the conductance value ( $83 \text{ S cm}^2 \text{ mol}^{-1}$ ) is an indication of its 1 : 1 electrolytic nature<sup>12</sup>.



TABLE-I  
ANALYTICAL AND MOLAR CONDUCTANCE DATA OF METAL COMPLEXES

Complex	Analysis, % Found (Calcd.)						Molar conductance ( $\text{S cm}^2 \text{ mol}^{-1}$ )
	M	C	H	N	S	Cl	
Ligand	—	60.53 (60.50)	4.95 (5.02)	11.76 (11.76)	8.96 (8.90)	—	—
[CoL <sub>2</sub> Cl <sub>2</sub> ]	6.94 (6.98)	51.10 (51.17)	4.53 (4.53)	9.92 (9.94)	7.43 (7.58)	8.13 (8.41)	4.79
[NiL <sub>2</sub> Cl <sub>2</sub> ].2H <sub>2</sub> O	6.63 (6.57)	49.00 (49.09)	4.30 (4.35)	9.58 (9.55)	7.20 (7.27)	8.00 (8.06)	8.78
[CuLCl(H <sub>2</sub> O) <sub>2</sub> ]	13.00 (12.91)	43.49 (43.88)	4.97 (4.71)	8.62 (8.53)	6.47 (6.50)	7.06 (7.20)	3.93
AgL(NO <sub>3</sub> )	20.32 (20.47)	40.51 (40.99)	3.44 (3.63)	10.45 (10.62)	6.40 (6.07)	—	0.46
[ZnL <sub>2</sub> SO <sub>4</sub> ]	7.70 (7.77)	49.64 (49.33)	4.41 (4.37)	9.38 (9.59)	11.03 (10.96)	—	6.70
[CdLCl <sub>2</sub> ]	20.60 (20.80)	39.68 (39.95)	3.56 (3.54)	7.66 (7.76)	5.90 (5.91)	13.19 (13.13)	8.64
[HgL <sub>2</sub> Cl <sub>2</sub> ]	20.46 (20.35)	43.91 (43.81)	3.82 (3.88)	8.59 (8.52)	6.44 (6.49)	7.17 (7.20)	10.18
[PbL <sub>2</sub> (NO <sub>3</sub> ) <sub>2</sub> ]	19.48 (19.82)	41.27 (41.32)	3.41 (3.66)	10.72 (10.71)	6.23 (6.12)	—	6.94
[FeL <sub>2</sub> Cl <sub>2</sub> ].Cl.5H <sub>2</sub> O	5.74 (5.78)	44.61 (44.69)	4.90 (4.97)	8.59 (8.69)	6.55 (6.62)	11.12 (11.02)	8.30

The cobalt complex [CoL<sub>2</sub>Cl<sub>2</sub>] shows a magnetic moment value of 4.50 BM. The higher than the spin only value for the magnetic moment is to be expected for Co(II) complex in tetrahedral environment<sup>13</sup>. The copper complex [CuLCl(H<sub>2</sub>O)<sub>2</sub>] is found to be diamagnetic, *i.e.*, complete reduction of Cu(II) → Cu(I) takes place during complex formation. It was observed during the synthesis of this complex that excess of ligand is required over the stoichiometric ratio and this proves that a part of the ligand which is in excess may reduce Cu(II) → Cu(I) and the cation thus formed complexes with the remaining ligand. This was confirmed by the synthesis of the same complex by prior reduction of Cu(II) salt by sulphur dioxide. For [NiL<sub>2</sub>Cl<sub>2</sub>].2H<sub>2</sub>O, the measured magnetic moment value is 3.73 BM which is higher than the spin-only value. Hence, orbital contribution may be expected from tetrahedral <sup>3</sup>T<sub>1</sub> ground state. For Fe(III) complex, the measured magnetic moment value 5.90 BM is close to spin-only

value for a  $d^5$  system. As expected, Zn(II), Cd(II), Hg(II), Pb(II) and Ag(I) complexes are diamagnetic.

The electronic spectrum of the ligand in DMF showed an absorption maximum at  $32,200\text{ cm}^{-1}$  ( $\epsilon = 190\text{ L mol}^{-1}\text{ cm}^{-1}$ ) due to  $n \rightarrow n^*$  transition and this remained almost unchanged in the complexes also. The electronic spectrum of Co(II) complex is dominated by a high energy intense band at  $18,900\text{--}16,500\text{ cm}^{-1}$ . This transition corresponds to  ${}^4A_2 \rightarrow {}^4T_1(P) = 12Dq + 15B$  ( $\nu_3$ ) and is responsible for the blue colour of the synthesized complex<sup>14</sup>. For a tetrahedral Co(II) complex two more transitions are theoretically possible,



Bands due to these transitions fall in IR and near IR regions and hence are not observed. By using the single transition energy, it is not possible to calculate crystal field parameters for Co(II) complex. However, the measured magnetic moment of Co(II) complex can be related to  $10Dq$ , by the relationship

$$\mu_{\text{eff}} = \mu_s(1 - 4\lambda/10Dq)$$

Substituting  $\mu_{\text{eff}} = 4.50\text{ BM}$ ,  $\mu_s = 3.87\text{ BM}$ ,  $\lambda = -180\text{ cm}^{-1}$ ,  $10Dq$  value of  $4425\text{ cm}^{-1}$  was estimated for Co(II) complex. This value falls in the range expected for tetrahedral Co(II) complexes and hence the assignment is apparently correct<sup>15</sup>.

For Ni(II) complex, there is an intense band at  $16,530\text{ cm}^{-1}$ . It is documented in literature<sup>16</sup> that tetrahedral Ni(II) complex shows high intensity bands in the region  $16,500\text{--}14,000\text{ cm}^{-1}$  and are attributed to  ${}^3T_1(F) \rightarrow {}^3T_1(P)$  transition ( $\nu_3$ ). A near IR band with appreciable intensity is also recorded at  $9600\text{ cm}^{-1}$ . This is assigned to  ${}^3T_1(F) \rightarrow {}^3A_2(F)$  transition. The ligand field parameters  $\nu_3/\nu_2 = 1.72$ ,  $B = 899\text{ cm}^{-1}$  and  $\beta = 0.85$  are quite consistent with tetrahedral geometry.

In case of Fe(III) complex, a number of weak spin forbidden bands are observed in the range  $20,400\text{--}18,000\text{ cm}^{-1}$ . These transitions are characteristic of tetrahedral Fe(III) complex<sup>13</sup> and hence tetrahedral geometry is proposed for the Fe(III) complex.

Thermogravimetric study of the Ni(II) and Fe(III) complexes has been carried out. In case of Ni(II) complex, a weight loss of 4.00% (theoretical 4.09%) in the temperature below  $150^\circ\text{C}$ <sup>16,17</sup> indicates the presence of two molecules of water of crystallization. This was followed by a single step mass-loss in the temperature range of  $200\text{--}500^\circ\text{C}$ , corresponding to all organic matter. The mass of the solid residue obtained was 8.40% of the initial mass, which corresponds to NiO (theoretical = 8.49%). While in Fe(III) complex, up to  $150^\circ\text{C}$ , the thermogram showed 9.30% weight loss (theoretical 9.31%) and this corresponds to five molecules of water of crystallization<sup>16,17</sup>. The complex showed maximum mass loss in the temperature range  $200\text{--}500^\circ\text{C}$ , corresponding to the loss of all organic matter. Final mass of residue in this case corresponds to  $\text{Fe}_2\text{O}_3$  (16.00%, theoretical = 16.52%).

In the complexes,  $^1\text{H}$  NMR signals, especially the singlet due to  $-\text{N}-\text{CH}_2-\text{N}-$  (compared to ligand spectrum) shifts downfield, indicating the involvement of oxadiazole moiety in complexation.

In the complexes of Co(II), Ni(II), Cu(I), Fe(III), Zn(II) and Pb(II), the shifts in band positions, compared to those present in the ligand molecule, are similar, thereby indicating the presence of similar type of bonding. The  $\nu(\text{C}-\text{O}-\text{C})$  of oxadiazole ring shifts downward by  $10-20\text{ cm}^{-1}$  in complexes, indicating the involvement of oxadiazole ring in complexation<sup>18</sup>. The  $\nu(\text{C}=\text{N})$  of these complexes shifts downward by about  $20\text{ cm}^{-1}$ , suggesting the bonding to metal through N-atom at 4-position. Though oxadiazole ring has two nitrogen atoms, at positions 3 and 4, it is almost likely that the one at 4-position is involved in bonding by taking into consideration the steric factors. In the case of Hg(II) and Ag(I) complexes, the  $\nu(\text{C}-\text{N})$  almost remains unaffected from its position in the ligand. But the  $\nu(\text{C}=\text{S})$  shows a significant downward shift (*ca.*  $15\text{ cm}^{-1}$ ), thereby indicating that the bonding might have taken place through the thione S atom<sup>11</sup>.

But in the Cd(II) complex, shifts in both  $\nu(\text{C}=\text{N})$  and  $\nu(\text{C}=\text{S})$  were observed. This means that in this case both the N-atom are involved, with the oxadiazole moiety acting as either a bidentate ligand or a bidentate bridging ligand. The most likely mode of bonding could be as a bidentate bridging ligand.

A broad bands in the range  $3500-3450\text{ cm}^{-1}$  in Ni(II), Fe(III) and Cu(I) complexes are due to  $\nu(\text{OH})$ <sup>19</sup>. This gives evidence for water of crystallization in Ni(II) and Fe(III) complexes. In the Cu(I) complex, a band at  $872\text{ cm}^{-1}$  supports the existence of coordinated water molecule<sup>20</sup>. In Ag(I) and Pb(II) complexes, bands at  $1421\text{ cm}^{-1}$   $\nu_{\text{asym}}(\text{NO}_2)$ ,  $1310\text{ cm}^{-1}$   $\nu_{\text{sym}}(\text{NO}_2)$  and  $1047\text{ cm}^{-1}$   $\nu(\text{NO})$  are due to monodentate nitrate group<sup>21</sup>. Bands at  $1105$  and  $1032\text{ cm}^{-1}$  are due to chelating sulphato group in Zn(II) complex<sup>22</sup>. The low frequency bands in the  $465-455$ ,  $445-435$  and  $380-370\text{ cm}^{-1}$  region are tentatively assigned to  $\nu(\text{M}-\text{N})$ ,  $\nu(\text{M}-\text{O})$  and  $\nu(\text{M}-\text{S})$ , respectively.

Antibacterial activity shown by the ligand and the complexes is given in Table-2. Concentration of compound is  $5\text{ mg}/10\text{ mL}$  in DMF. Organisms studied are *Bacillus subtilis*, *Staphylococcus aureus*, *Escherichia coli* and *Pseudomonas aeruginosa*.

TABLE-2  
RESULTS OF ANTIBACTERIAL ACTIVITY SCREENING  
OF LIGAND AND METAL COMPLEX

Compound	Diameter of zone of inhibition (mm)			
	<i>B. subtilis</i>	<i>S. aureus</i>	<i>E. coli</i>	<i>Ps. aeruginosa</i>
Ligand	12	18	30	12
$[\text{FeL}_2\text{Cl}_2]\text{Cl}\cdot 5\text{H}_2\text{O}$	18	20	48	24
$[\text{CoL}_2\text{Cl}_2]$	12	20	28	12
$[\text{NiL}_2\text{Cl}_2]\cdot 2\text{H}_2\text{O}$	14	18	28	14
$[\text{CuLCl}(\text{H}_2\text{O})_2]$	12	19	29	12
$[\text{ZnL}_2\text{SO}_4]$	14	20	28	12

Our results of investigation on the antibacterial activity revealed that the Fe(III) complex exhibits much higher activity towards all the organisms studied than the

ligand. The results indicate that the Fe(III) complex showed enhanced antibacterial activity towards *B. subtilis*, *E. coli* and *Ps. aeruginosa*. The other metal complexes did not show any significant increase in activity as compared to the ligand.

### Conclusion

The ligand molecule acts as a monodentate ligand in all the studied cases except in Cd(II) complex, bonding either through N(4) or through S—depending upon the nature of the metal ions. In the Cd(II) complex, it appears to act as a bidentate bridging ligand. Tetrahedral structures for Fe(III), Co(II), Ni(II), Zn(II), Cu(I), Hg(II) and Pb(II) complexes, tetrahedral polymeric structure for Cd(II) and linear structure for Ag(I) complex have been tentatively proposed.

### ACKNOWLEDGEMENTS

The author is thankful to Heads of RSIC, IIT Delhi and RSIC, CDRI Lucknow for recording the spectra and for elemental analyses.

### REFERENCES

1. T. Ramalingam, A.A. Desmukh and P.B. Sattur, *J. Indian Chem. Soc.*, **58**, 269 (1981).
2. N. Soni, J.P. Barthwal, T.K. Gupta, T.N. Bhalla and K.P. Bhargava, *Indian Drugs*, **19**, 301 (1982).
3. B. Kalluraya, R. Chimbalkar, P. Gunaya, *Indian J. Heterocycl. Chem.*, **6**, 103 (1996).
4. P.R. Kagthara, N.S. Shah, R.K. Doshi and H.H. Parekh, *Indian J. Chem.*, **38B**, 572 (1999).
5. A. Kocabalkanli, O. Ates and G. Otuk, *Il Farmaco*, **56**, 975 (2001).
6. F. Bentiss, M. Traisnel and M. Lagrenee, *Corrosion Sci.*, **42**, 127 (2000).
7. R.A. Rai, U. Agarwala and Lakshmi, *Indian J. Chem.*, **18A**, 426 (1979).
8. R.W. Young and K.H. Wood, *J. Am. Chem. Soc.*, **77**, 400 (1955).
9. B. Kalluraya, R. Chimbalkar and B.S. Holla, *Indian J. Heterocycl. Chem.*, **5**, 37 (1995).
10. G.H. Jeffery, J. Basett, J. Mendham and R.C. Denny (Eds.) A.I. Vogel's Text Book of Quantitative Inorganic Analysis, 5th Edn., Addison-Wesley-Longman Inc. p. 326 (1989).
11. Alan R. Katritzky and Charles W. Rees, *Comprehensive Heterocyclic Chemistry*, Pergamon Press, Vol. 6, p. 25 (1984).
12. W.J. Geary, *Coord. Chem. Rev.*, **7**, 81 (1971).
13. B.N. Figgis and J. Lewis, *Prog. Inorg. Chem.*, **6**, 185 (1964).
14. F.A. Cotton, B.M.L. Goodgame and M. Goodgame, *J. Am. Chem. Soc.*, **83**, 4690 (1961).
15. R.S. Drago, *Physical Methods for Chemists*, Saunders College Publishing, p. 448 (1992).
16. K.C. Satpathy, R. Mishra and B.B. Jal, *J. Indian Chem. Soc.*, **63**, 377 (1986).
17. V. Mishra and M.C. Jain, *J. Indian Chem. Soc.*, **65**, 28 (1988).
18. R. Nigam, S. Swarup, V.K. Saxena and H.K. Singh, *J. Indian Chem. Soc.*, **69**, 692 (1992).
19. F.A. Miller and C.H. Wilkins, *Anal. Chem.*, **24**, 1253 (1952).
20. I. Nakagawa and T. Shimanouchi, *Spectrochim. Acta*, **20**, 429 (1964).
21. B.M. Gatehouse, S.E. Livingstone and R.S. Nyholm, *J. Chem. Soc.*, 4222 (1957); *J. Inorg. Nucl. Chem.*, **8**, 75 (1958).
22. C.G. Barraclough and M.L. Tobe, *J. Chem. Soc.*, 1993 (1961).