

Spectrophotometric Characterisation of Doped Pr(III) Ion in Saturated Alcoholic Solution of Some Thiosemicarbazone derived from Carbonyl Compounds

ANU ARORA, ANJU POPLI, P.C. TATER
and K.P. SONI*

*Department of Chemistry
Government Dungar College
Bikaner-334 001, India*


The stereo-environment of doped Pr(III) ion in saturated solution of some thiosemicarbazones of carbonyl compounds in alcoholic media has been discussed on the basis of various electronic spectral parameters. The various electronic parameters, viz., Judd-Ofelt intensity parameter, (T_{λ}) Slater-Condon (F_k), Lande's (ζ_{4f}) intensity of hypersensitive band (3P_2) and Racah's (E_k) as well as bonding parameter $b^{1/2}$ for Pr(III) ion doped in alcoholic saturated solution of thiosemicarbazones derived from thiosemicarbazide and ethyl vanillin, 3,4-dimethoxy benzaldehyde, 2,5-dimethoxy benzaldehyde, chalcone, *p*-hydroxy acetophenone, benzophenone, 3-nitrobenzaldehyde and 4-amino acetophenone etc.

INTRODUCTION


In the present discussion the study has been made on the basis of utility of some lanthanide substituted thiosemicarbazones of carbonyl compounds, Owing to their immense medicinal application¹⁻³, on the basis of Judd-Ofelt and Slater-Condon theory. Since in the field of science and technology the doped study of certain systems finds immense application⁴⁻¹³, therefore, on the basis of Judd-Ofelt¹⁴⁻¹⁷ intensity as well as other electronic spectral parameters a correlation of $f \leftrightarrow f$ transition for Pr(III) ion doped in saturated solution of some thiosemicarbazones of carbonyl compounds and the stereo-environment around Pr(III) ion has been attempted. A saturated solution in alcoholic media has been made by dissolving some thiosemicarbazones derived from carbonyl compounds at room temperature ($30 \pm 2^\circ\text{C}$) and a constant amount of Pr(III) chloride has been added to each of the solutions. Characterisation of some of the electronic parameters for Pr(III) ion doped in saturated alcoholic solution of these compounds was made spectrophotometrically. On the basis that Pr(III) ion yields four bands in visible region, viz., 3P_2 , 3P_1 , 3P_0 and 1D_2 , the change in the intensity of these bands is indicated by red shift caused by the change in coordination environment around the ion resulting from $f \leftrightarrow f$ transition in lanthanides.

EXPERIMENTAL


Equal molar concentration of thiosemicarbazide and ethyl vanillin, 3,4-dimethoxy benzaldehyde, 2,5-dimethoxy benzaldehyde, chalcone, *p*-hydroxy acetophenone, benzophenone, *m*-nitrobenzaldehyde and *p*-amino acetophenone were refluxed for 5 to 6 h at temperature 70 to 80°C; thiosemicarbazones derived as such were crystallized and various saturated solutions¹⁸⁻²³ of above compounds and 0.014 M of PrCl₃·6H₂O (99.9% purity supplied by Indian Rare Earth's Udyogmandalam, Kerala) were added to each of the solutions. The solution spectra of these systems were recorded by using standard spectrophotometer²⁴⁻²⁹ (Spectronic-20) in the visible region. The calculation for various electronic parameters was made by computerized statistical method reported earlier²⁶⁻²⁹.




Ethyl vaniline thiosemicarbazone




3,4-Dimethoxy benzaldehyde thiosemicarbazone




2,5-Dimethoxy benzaldehyde thiosemicarbazone




Chalcone thiosemicarbazone




Benzophenone thiosemicarbazone




4-Hydroxy acetophenone thiosemicarbazone



Phenazone thiosemicarbazone



3-Nitro benzaldehyde thiosemicarbazone



4-Amino acetophenone thiosemicarbazone

RESULTS AND DISCUSSION

The analytical data of thiosemicarbazones are given in Table-1. The absorption at 1600 cm^{-1} in the thiosemicarbazones can be attributed to $\nu(\text{C}=\text{N})$ of imine nitrogen which is in agreement with early reports^{18,19}. The bands observed in $1630\text{--}1130\text{ cm}^{-1}$ region are assigned to $\nu(\text{C}=\text{S})$ ²⁰⁻²¹. The computed values of various parameters have been given in Tables 2 and 3. The present study shows that the $f \leftrightarrow f$ transition resulting from spin orbit interaction in Pr(III) is due to interaction with the ligand present in the saturated solution which is remarkably manifested in terms of change in the energy of various electronic parameters. The decrease in computed values of spin orbit interaction parameter (ζ_{4f}), oscillator strength of ${}^3\text{P}_2$ band, Judd-Ofelt parameter (T_λ) and Slater-Condon parameter (F_k) in comparison to free ion values clearly reflects symmetry change around the metal ion in the solution. The T_4/T_6 ratio reflects the change in stereo-environment around the metal ion present in the solution and thereby change in coordination number. Therefore the change in the surrounding environment around the metal ion involving the suitable interaction between the metal ion and

ligand has been well characterized by decrease in the various inter-electronic repulsions and spin-orbit interactions.

TABLE-1
ANALYTICAL DATA OF THIOSEMICARBAZONES

Thiosemi-carbazones	% Analysis, Found (Calcd.)			
	C	H	N	S
1	52.17 (52.20)	5.92 (5.90)	16.60 (16.61)	12.64 (12.60)
2	50.20 (50.18)	5.43 (5.40)	17.57 (17.55)	13.38 (13.40)
3	50.20 (50.22)	5.43 (5.40)	17.57 (17.60)	13.38 (13.35)
4	68.57 (68.60)	5.00 (4.49)	15.00 (15.04)	11.42 (11.31)
5	65.88 (65.42)	5.09 (4.48)	16.47 (16.50)	12.54 (12.44)
6	51.67 (51.68)	5.26 (5.25)	20.09 (20.05)	15.31 (15.32)
7	54.75 (54.80)	6.46 (6.52)	26.61 (25.92)	12.16 (12.20)
8	42.85 (42.70)	3.57 (3.62)	25.0 (24.95)	14.28 (14.11)
9	51.92 (50.99)	5.76 (5.62)	26.92 (27.13)	15.38 (15.43)

TABLE-2
COMPUTED VALUES OF ENERGY (cm^{-1}) OF OSCILLATOR STRENGTH T_λ AND T_4/T_6 FOR Pr(III) ION DOPED IN ALCOHOLIC SATURATED SOLUTION OF VARIOUS THIOSEMICARBAZONES OF CARBONYL COMPOUNDS

Com-pounds	${}^3P_2 \times 10^{-6}$		${}^3P_1 \times 10^{-6}$		${}^3P_0 \times 10^{-6}$		${}^1D_2 \times 10^{-6}$		$T \times 10^{-10}$			T_4/T_6
	P_{exp}	P_{cal}	P_{exp}	P_{cal}	P_{exp}	P_{cal}	P_{exp}	P_{cal}	T_2	T_4	T_6	
1	24.0	24.0	9.67	9.67	3.50	3.49	3.61	3.61	-778	18.1	74.1	0.24
2	19.8	19.8	7.73	7.73	2.91	2.90	3.81	3.81	-450	14.5	61.1	0.24
3	27.9	27.9	11.40	11.40	4.87	4.85	4.24	4.24	-879	22.4	84.9	0.26
4	28.5	28.5	11.70	11.70	5.08	5.06	5.17	5.17	-713	23.1	86.9	0.27
5	20.6	20.6	8.05	8.05	3.05	3.04	3.11	3.11	-659	15.2	63.4	0.24
6	24.6	24.6	12.80	12.80	4.36	4.34	4.72	4.72	-569	23.6	74.4	0.32
7	28.7	28.7	11.80	11.80	4.99	4.96	4.31	4.31	-937	23.1	88.3	0.26
8	27.7	27.7	10.07	10.07	3.97	3.95	5.14	5.14	-676	20.2	85.7	0.24
9	23.2	23.2	7.37	7.37	2.74	2.72	2.85	2.85	-891	13.9	72.4	0.19

TABLE-3
 COMPUTED VALUES OF F_k , ζ_{4f} , E_k , $\%F_2$, β and $b^{1/2}$ Pr(III) ION DOPED IN ALCOHOLIC SATURATED SOLUTIONS OF VARIOUS THIOSEMICARBAZONES OF CARBONYL COMPOUNDS

Compounds	F_2	F_4	F_6	$\%rF_2$	ζ_{4f}	$\%r\zeta_{4f}$	E^1	E^2	E^3	β	$b^{1/2}$
Pr ³⁺ ion	322.09	44.46	4.87	-	738.00	-	4729.00	24.78	478.13	-	-
Compound No. 1	311.39	42.98	4.70	3.31	672.69	9.21	4571.86	23.91	462.23	0.96680	0.12884
Compound No. 2	311.03	42.93	4.69	3.43	692.31	6.56	4566.50	23.88	461.69	0.96566	0.13102
Compound No. 3	310.71	42.89	4.69	3.53	694.46	6.27	4561.83	23.86	461.22	0.96467	0.13289
Compound No. 4	312.81	43.18	4.72	2.87	661.77	10.69	4592.72	24.02	464.34	0.97121	0.11997
Compound No. 5	310.60	42.87	4.69	3.56	701.23	5.36	4560.24	23.85	461.06	0.96434	0.13352
Compound No. 6	311.37	42.98	4.70	3.32	673.07	9.16	4571.51	23.91	462.20	0.96672	0.12898
Compound No. 7	308.45	42.58	4.66	4.23	715.66	3.41	4528.63	23.68	457.86	0.95765	0.14550
Compound No. 8	311.57	43.01	4.70	3.26	666.50	10.05	4574.48	23.92	462.50	0.96735	0.12776
Compound No. 9	308.06	42.52	4.65	4.35	731.29	1.30	4522.94	23.65	457.29	0.95645	0.14755

The parametric values provide useful information regarding the coordination behaviour of Pr(III) ion. The coordination behaviour varies as the change in ratio of T_4/T_6 takes place. The similar values indicate the similar stereo-environment around Pr(III) due to solute-interaction present in the solution.

The values of Slater-Condon parameter (F_k) in the present system decrease from free ion value and show close similarity whereas Lande's parameter (ζ_{4f}) which shows spin-orbit interaction has been found to show much variation. This shows that spin-orbit interactions in $f \leftrightarrow f$ transition are much more affected in comparison to inter-electronic interaction. The intensity of hypersensitivity band also shows similar trend.

The study provides useful information regarding stereo-environment around Pr(III) ion as significantly exhibited by T_4/T_6 ratio. The nephelauxetic ratio as indicated by the values of $b^{1/2}$ also reflects some bonding character in the form of change in stereo-environment around the central metal ion.

The various other electronic parameters reported in Tables 2 and 3 have their usual significance as reported earlier by Soni *et al.*^{31, 32}.

ACKNOWLEDGEMENTS

Authors are thankful to Prof. S.L. Jain and H.K. Pandey, Anil Gupta, Narendra Bhojak, Department of Chemistry, Dungar College, Bikaner for their valuable suggestions and help. One of the authors K.P. Soni is also thankful to UGC for financial assistance.

REFERENCES

1. David B. Chapman and E. Leongway, *Ann. Rev. Pharmacol. Toxicol.*, **553** (1980).
2. B. Weiss George, *Cellular Pharmacology of Lanthanum*, Vol. 14, p. 343 (1974).
3. S.I. Levy, *The Rare Earths*, Longmans Green, New York, p. 171 (1915).
4. B.R. Judd, *Phys. Rev.*, **127**, 750 (1962).
5. ———, *J. Chem. Phys.*, **44**, 839 (1966).
6. W.T. Carnall, P.R. Fields and K.J. Rajank, *Chem. Phys.*, **49**, 4412, 4424, 4443, 4447, 4450 (1968).
7. G.S. Ofelt, *J. Chem. Phys.*, **37**, 511 (1962).
8. W.T. Carnall and P.R. Field, *Chem. Phys.*, **42**, 3797 (1965).
9. S.P. Tandon and P.C. Mehta, *J. Chem. Phys.*, **52**, 4313 (1970).
10. R.D. Peacock, *J. Chem. Soc. (A)*, 2028 (1971).
11. S.P. Tandon and P.P. Vaishnava, *Can. J. Spectra*, **22**, 17 (1970).
12. S.P. Tandon and P.C. Mehta, *J. Chem. Phys.*, **52**, 4313, 4896 (1970).
13. S.P. Tandon, P.P. Vaishnava and R.C. Mathur, *Sol. State Comm.*, **15**, 787 (1974).
14. S.N. Mishra, G.K. Joshi and Megh Singh, *J. Inorg. Nucl. Chem.*, **43**, 206 (1981).
15. S.N. Mishra and G.K. Joshi, *Indian J. Pure Appl. Phys.*, **19**, 279 (1981).
16. S.N. Mishra, Surendra B. Mehta and Venkat K. Subramaniyan, *Asian J. Chem. Rev.*, **2**, 100 (1991).
17. S.P. Tandon and R.C. Govil, *Z. Naturforsch.*, **26(a)**, 1357 (1971).
18. P.S. Radha Krishnan, P. Indrasenan and C.G.R. Nair, *Polyhedron*, **3**, 67 (1984).
19. R.K. Agarwal and Indranil Chakravarti, *Synth. React. Inorg. Met.-Org. Chem.*, **25**, 679 (1995).

20. K. Swaminathan and H.M.N.H. Irving, *J. Inorg. Nucl. Chem.*, **26**, 1291 (1964).
21. D. Banerjee and I.P. Singh, *Indian J. Chem.*, **6**, 34 (1968).
22. G.K. Joshi, Megh Singh and S.N. Mishra, *Indian J. Chem.*, **23A**, 329 (1984).
23. G.K. Joshi, M.P. Bhutra and S.N. Mishra, *J. Inorg. Nucl. Chem.*, **43**, 525 (1981).
24. G.K. Joshi, *Indian J. Pure and Appl. Phys.*, **21**, 224 (1983).
25. G.S. Manku, *Theoretical Principles of Inorganic Chemistry*, Tata McGraw-Hill, New Delhi, p. 60 (1980, reprint 1992).
26. S.V.J. Lakshman and Janardhanam, *Proceedings of the National Symposium of Optical and ESR Spectra of Transition Metal and Rare Earth Ions*, Srivenkateshwara University, Tirupati, pp. 18–19 (11–15 Jan. 1982).
27. S.P. Tandon, A. Gokhroo, M.P. Bhutra, S.S.L. Surana and K. Tandon, *J. Chem. Phys.*, **81**, 4243 (1987).
28. S.S.L. Surana, Megh Singh and S.N. Mishra, *J. Inorg. Nucl. Chem.*, **42**, 61 (1980).
29. P.C. Vyas, C.K. Ojha, Ms. S. Mittal and G.K. Joshi, *J. Indian Chem. Soc.*, **65**, 611 (1988).
30. B.K. Gupta, G.K. Joshi and P.R. Bhati, *Indian J. Applied Physics*, **28**, 525 (1990).
31. P.R. Bhati, K.P. Soni, G.K. Joshi and S.N. Swami, *Asian J. Chem.*, **4**, 828 (1992).
32. P.R. Bhati, K.P. Soni, P.C. Tater, G.K. Joshi and S.N. Swami, *Asian J. Chem.*, **5**, 857 (1993).
33. (Miss) Daya Balani, (Mrs.) Shakuntala Maurya, G.K. Joshi and P.C. Tater, *Asian J. Chem.*, **6**, 801 (1994).

(Received: 7 August 2000; Accepted: 1 November 2000)

AJC-2143