



Accumulation of Copper and Lead in Selected Medicinal Plants

S. SHAVAKH¹, S. GHALANDARI KAVKANI^{1,*} and J. NOURI²

¹Department of Plant Protection, School Faculty of Agriculture and Natural Resources, Science and Research Branch, Islamic Azad University, Tehran, Iran

²Department of Environmental Management, Graduate School of the Environment and Energy, Science and Research Branch, Islamic Azad University, Tehran, Iran

*Corresponding author: E-mail: s_ghalandary@yahoo.com

(Received: 16 January 2012;

Accepted: 5 December 2012)

AJC-12500

The present study aims to estimate lead and copper metals content in four medicinal plants collected from different sites of Saveh region located in south western of Tehran, in June and August 2010. Samples of *Eucalyptus globules*, *Mentha piperita*, *Thymus vulgaris* and *Rosa damascena* were separately collected from different locations and prepared in laboratory. Lead and copper content were measured in each samples using atomic absorption spectrophotometry. Results show that lead content was the highest in *Mentha piperita* and the lowest was in *T. vulgaris*. However, *T. vulgaris* and *E. globulus* contained the highest and the lowest levels of copper, respectively. Overall, despite of vital advantages of studied medicinal plants, heavy metals contamination prevent of therapeutic uses of these plants which may result in fatal effects on human health.

Key Words: Metal pollution, *Eucalyptus globules*, *Mentha piperita*, *Thymus vulgaris*, *Rosa damascena*.

INTRODUCTION

During past decades, the use of medicinal plants in agronomy productions, dietary supplements, pharmacy, as well as therapeutic purposes has been achieved with increasing interest worldwide¹⁻⁴. In this regard, herbal medicine has also played a crucial role in the pharmaceutical and health markets of 21st century⁵. So that, growing interest of developing, as well as developed countries towards cultivation of medicinal and aromatic plants has augmented rapidly^{6,7}. However, there have been also reported some cases of illness and fatalities induced by use of such plants^{8,9}. Poisoning of toxic metals, contained in medicinal plants through soil, water or air, were reported in the United States, Europe and Asian countries, as well¹⁰⁻¹³. In addition, there exist some major reasons for presence of toxic metals in medicinal plants, including growth in polluted areas, use of agricultural expedients (such as cadmium-containing fertilizers, organic mercury or lead based pesticides); contaminated irrigation water¹⁴⁻¹⁷. Accumulation of metals can occur in both root and above-ground organs deduced to transfer toxic metals into food chain¹⁸. So, several harmful effects on animal and human health, such as nephrotoxicity which induce abnormalities of tubular re-absorption are as the consequences¹⁹. Moreover, renal and nervous system would be attacked by the adverse effects of lead and

mercury^{20,21}. For instance, copper facilitates Glucose oxidation and energy release consequently. Iron absorption, oxygen supplement for tissues particularly brain tissues and brain function improve by higher level of copper in body. Copper level is relatively related to estrogen level. Shortage of copper deduced to anemia, depression and dermal problems. However, high adsorption of copper leads to headache and hypoglycemia. Moreover, additional copper content is deposited in brain and liver which damage livers and result in anemia and alopecia in women. Besides, lead is a naturally-occurring element which might be harmful to humans, especially children, if ingested or inhaled higher. Lead poisoning can cause a number of adverse human health effects, but is particularly detrimental to the neurological development of children. In this regard, monitoring of heavy metals contained in medicinal plants is of great priority²² for two essential reasons: first, increasing trend of general environment contamination by toxic metals²³, second, toxic contains of heavy metals and/or arsenic content in exotic herbal remedies. The present study aims to determine the amounts of copper and lead in four selected medicinal plants, including *Eucalyptus globules*, *Mentha piperita*, *Thymus vulgaris* and *Rosa damascena*. This research has been performed in Science and Research Branch, Islamic Azad University, Tehran, during 2010-2011.

EXPERIMENTAL

Sampling of *Eucalyptus globules*, *Mentha piperita*, *Thymus vulgaris* and *Rosa damascena* were performed randomly in June and August 2010 throughout the Saveh region located in south western of Tehran, Iran (Fig. 1). The samples were firstly rinsed with distilled deionized water and then air dried in the laboratory condition. Then, each sample was milled separately in a micro-hammer (without metal in it) and 2 g of powdered sampled plants were poured in Erlenmeyer adding 20 mL HNO₃. Digesting the composed samples on a hot plate exposure to a low temperature (40 °C), 20 mL HNO₃ 2 N has been added and the total volume reached to 50 mL. The sample was then vacuum-filtered through Whatman 42 filter paper. Afterwards, by Atomic absorption spectrophotometry 200; varian; made in USA. Accumulation level of Cu and Pb were measured and shown in standard curves and graphs.



Fig. 1. Location of the study area

Table-1 shows the studied medicinal plants specifying therapeutically usable parts of them for the present study.


TABLE-1 SCIENTIFIC TITLES OF SAMPLED PLANTS SPECIFYING THERAPEUTICALLY USABLE PARTS		
Sample code	Family	Scientific name
a	Myrtaceae	<i>Eucalyptus globulus</i>
b	Lamiaceae	<i>Mentha piperita</i>
c	Lamiaceae	<i>Thymus vulgaris</i>
d	Rosaceae	<i>Rosa damascena</i>

RESULTS AND DISCUSSION


Heavy metals concentrations were investigated in four selected medicinal plants and results are shown in Figs. 2 and 3. analyzing two elements (Cu and Pb), metal contents were in order of below: **Pb content:** *Mentha piperita* > *Eucalyptus globulus* > *Rosa damascena* > *Thymus vulgaris*. **Cu content:** *Thymus vulgaris* > *Rosa damascena* > *Mentha piperita* > *Eucalyptus globulus*, confirmed that the heavy metals contents in plants depending on the country of origin, environmental pollution levels, plant part and technological processes.

Lead: Lead is the most toxic environmental pollutant the main sources of lead pollution in agriculture and plants are lead mines, fuel combustion, sewage sludge applications and farmyard manure²⁴. It reacts or complexes with many biomolecules and adversely affects the reproductive, nervous,


gastrointestinal, immune, renal, cardiovascular, skeletal and muscular systems as well as developmental processes²⁵ suggest that lead levels are generally higher in leaves than in other plant parts probably because lead contaminations occur mainly through the atmosphere whereas lead uptake capacity of plants is generally low²⁶.




(a)
Elements



(b)
Elements



(c)
Elements



(d)
Elements

Fig. 2. Pb and Cu values in selected plants in 1st step




Fig. 3. Pb and Cu values in selected plants in 2nd step

Copper: Copper plays a role in the oxidative defense system, on the other hand chronic cu toxicity can results in severe poisoning, oxidative stress and human health²⁷. Copper is considered as a micronutrient for plants²⁸, Cu is also an essential component of various proteins like plastocyanin of photosynthetic system and cytochrome oxidase of respiratory electron transport chain²⁹. But enhanced industrial and mining activities have contributed to the increasing occurrence of Cu in ecosystems. Although copper is an essential enzymatic element for normal plant growth and development but can be toxic at excessive levels. Phytotoxicity can occur if its concen-

tration in plants is higher than 20 mg/kg DW (dry weight). Critical concentration for copper in plants is 20-100 mg/kg³⁰. Overall, therapeutic properties of the two latter plants are of high importance, due to Cu content as a harmful element for human health, they are not suggested for consumption.

Throughout the conducted investigation, copper and lead level were measured consecutively and tabulated in Tables 2 and 3. According to the results, lead level in *Eucalyptus globulus*, as well as *Mentha piperita* was higher than copper. However, *Thymus vulgaris* and *Rosa damascena* contained more copper rather than lead.

Sample	Conc. (mg/L)	RSD (%)
Phase 1		
a	0.290	2.2
b	0.400	1.1
c	0.080	36.5
d	0.020	>100
Phase 2		
a	0.310	-
b	0.404	-
c	0.082	-
d	0.021	-

Sample	Conc. (mg/L)	RSD (%)
Phase 1		
a	0.190	0.6
b	0.260	0.8
c	0.286	0.3
d	0.230	0.9
Phase 2		
a	0.195	-
b	0.263	-
c	0.290	-
d	0.240	-

RSD: Relative standard deviation.

Conclusion

Medicinal plants gained an important role in different pharmaceutical as well as therapeutic purposes during last decades. However, accumulation of heavy metals in these plants induces more attention before using them for remedies diseases. Therefore, it is essential to monitor the heavy metals content if such plants are meant for human consumption. Further studies are recommended for heavy metal contents of medicinal plants. The content of toxic metals in plant spices and herbs was found to be generally low. The highest contents of cooper was shown in *Eucalyptus globules*, the highest contents of lead was shown in *Mentha piperita*.

ACKNOWLEDGEMENTS

The authors thank Mr. A.R. Soltanabadi and S. Salehi for useful corporation throughout the progress of this work and the Science and Research Branch, the Laboratories Complex for its support throughout the analytical experiments.

REFERENCES

1. P.W. Wood, *Essence*, **30**, 42 (1999).
2. I.A. Khan, J. Allgood, L.A. Walker, E.A. Abourashed, D. Schelenk and W.H. Benson, *J. AOAC Int.*, **84**, 936 (2001).
3. WHO, Drug Information, Herbal Medicines, World Health Organization, Geneva, Vol. 16 (2002).
4. H. Krug, Gemuseproduktion Paul Parey, Cited from Chizzola, R. Metallic Trace Elements in Herbs and Spices Grown in Austria *Acta Hort.*, p. 249 (1986).
5. H.E. Kleinschmidt and R.W. Johnson, Weeds of Queensland, Queensland Department of Primary Ind., Australia, p. 147 (1977).
6. E. Liu and Y.J. Zheng, *Asian J. Chem.*, **23**, 1091 (2011).
7. J. Nouri, B. Lorestani, N. Yousefi, N. Khorasani, A.H. Hasani, F. Seif and M. Cheraghi, *Environ. Earth Sci.*, **62**, 639 (2011).
8. M.J. Stewart, J.J. Moar, P. Steenkamp and M. Steenkamp, *Forensic Sci. Int.*, **101**, 177 (1999).
9. E. Ernst, *Pharmacol. Sci.*, **23**, 136 (2002).
10. A. Olujuhunbe, P.A. Fields and A.F. Sanford, *Postgraduate Med. J.*, **70**, 764 (1994).
11. D.W. Dunbabin, G.A. Tallis and P.Y. Popplewell, *Med. J. Aust.*, **157**, 835 (1992).
12. T. Kaksoy, A. Hudak and M. Naray, *J. Toxicol.-Clin. Toxicol.*, **34**, 507 (1996).
13. S.B. Markowitz, C.M. Nenez and J.M. Clark, *Fields Crops Res.*, **60**, 143 (1999).
14. C.M.D. Levitt, *J. Am. Med. Assoc.*, **252**, 3127 (1984).
15. T.Y.K. Chan, B. Tomlinson and A.J.H. Critchley, *Lancet*, **342**, 1532 (1993).
16. E. Pip, *Bull. Environ. Contam. Toxicol.*, **46**, 790 (1991).
17. A.A.K. Abou-Arab, M.S. Kawther, M.E. El Tantawy, R.I. Badeaa and N. Khayria, *Food Chem.*, **67**, 357 (1999).
18. A. Kabata-Pendias and H. Pendias, Trace Elements in Soils and Plants, CRC Press Inc, Boca Raton, Florida, USA, edn. 2 (1992).
19. G. Nordberg, *Toxicol. Pharmacol.*, **30**, S57 (1999).
20. S. Tong, Y.E. von Schirnding and T. Prapamontol, *Bull. World Health Organization*, **78**, 1068 (2000).
21. WHO, Elemental Mercury and Inorganic Mercury Compounds: Human Health Aspects, Concise International Chemical Assessment Document 50, World Health Organization, Geneva (2003).
22. P.A.G.M. De Smet, Adverse Effects of Herbal Drugs; Springer-Verlag Berlin; ISBN 3-540-53100-9 (1992).
23. S.L. Ali, *Pharm. Ind.*, **45**, 1294 (1983).
24. M.A. Khan, I. Ahmad and Inayat-ur-Rahman, *J. Chin. Chem. Soc.*, **54**, 339 (2007).
25. F.M. Johnson, *Mut. Res.*, **410**, 123 (1998).
26. A.A. Abou-Arab and M.A.A. Donia, *J. Agric. Food Chem.*, **48**, 2300 (2000).
27. J.Y. Uriu-Adams and K.C.L. Copper, *Mol. Aspects Med.*, **26**, 268 (2005).
28. F. Thomas, C. Malick, E.C. Endreszl and K.S. Davies, *Physiol. Plant.*, **102**, 360 (1998).
29. K. Demirevska-Kepova, L. Simova-Stoilova, Z. Stoyanova, R. Holzer and U. Feller, *Environ. Exp. Botany*, **52**, 253 (2004).
30. U. Gupta, In ed.: J.O. Nariago, Copper in the Environment, John Wiley & Sons, New York, p. 255 (1975).