

Study of Tetragonally Compressed Complexes of Fe(II), Ni(II) and Cu(II)

SUDAMA CHAUDHRY¹, SUMIT KUMAR SINHA¹, PARVEEN KUMAR¹, PUNIT KUMAR¹, TRILOK PRASAD² and Shivadhar Sharma^{1,*}

¹Department of Chemistry, Magadh University, Bodh Gaya-824 234, India ²Department of Chemistry, Ram Nagina Singh College, Maner, Patna-801 111, India

*Corresponding author: Tel: +91 631 2220039; E-mail: sharma.shivadhar@gmail.com

Received: 23 June 2017;	Accepted: 27 September 2017;	Published online: 30 November 2017;	AJC-18652
-------------------------	------------------------------	-------------------------------------	-----------

4'-Nitrobenzaldehyde-4-phenylsemicarbazone and 4'-nitrobenzaldehyde-4-phenylthiosemicarbazone have been used as primary ligands for complexatation with Fe(II), Ni(II) and Cu(II). The complexes have been formulated as $[M(NBPS)_2X_2]$ and $[M(NBPTS)_2X_2]$ where, M = Fe(II), Ni(II) and Cu(II), NBPS = 4'-nitrobenzaldehyde-4-phenylsemicarbazone and NBPTS = 4'-nitrobenzaldehyde-4-phenyl thiosemicarbazone and X = CH₃COO⁻. The IR spectra of the metal-complexes in comparison to free ligands reveal that ligands act as monoanionic bidentate forming five membered chelate rings around the metal ion. The electronic spectra of the metal-complexes indicate tetragonally distorted octahedral geometry around the metal ion in complexes. The various crystal field parameters have been derived from the electronic spectra of the complexes. The greater value of Dq_(z) in comparison to the Dq_(xy) values as well as the negative values of D_t clearly indicate tetragonally compression along Z-axis. The complexes have also been evaluated for their biological activities and it has been observed that biological activities of free ligands enhanced after complex formation.

Keywords: Tetragonally compressed octahedral, Dq_(z), D_t.

INTRODUCTION

The degeneracy of orbitals can be removed by the lowering of symmetry of molecules. This can be achieved by Jahn-Teller distortion theorem. This may be stated as - "In a non-linear molecule if degenerate orbitals are asymmetrically occupied a distortion occurs to remove the degeneracy." or it may also be stated as "In an electronically degenerated state a non-linear molecule undergoes distortion to remove the degeneracy by lowering the symmetry and thus by lowering the energy" [1-3]. The distortion may also be caused by putting different ligands in XY-plane and along Z-axis as they are likely to produce crystal field of different strength in XY-plane and along Z-axis. The tetragonally distortion may either be Z-out distortion or Z-in distortion depending upon elongation along Z-axis or compression along Z-axis. The elongation along Zaxis is most preferred and has been widely studied in respect of Z-in distortion [4-8].

Semicarbazones and thiosemicarbazones have imerged as promising ligands in the coordination chemistry in the light of their excellent binding capability. The literature survey revealed a plethora of work in the field of semicarbazones and thiosemicarbazones and their transition metal complexes because of their fascinating structural properties and broad spectrum of their application in various field [9-15]. Due to their pharmaceutical properties which are frequently higher for the metal complexes than the free ligands, they have extensively been studied during recent years.

The present paper deals with the synthesis, structural and biological evaluation of tetragonally compressed octahedral complexes of 4'-nitrobenzaldehyde-4-phenylsemicarbazone and 4'-nitrobenzaldehyde-4-phenylthiosemicarbazone with Fe(II), Ni(II) and Cu(II).

EXPERIMENTAL

All the chemicals used were of AnalR Grade. 4'-Nitrobenzaldehyde was procured from Fluka and 4-phenylsemicarbazide and 4-phenylthiosemicarbazide were purchased from Sigma Aldrich. Fe(CH₃COO)₂, Ni(CH₃COO) and Cu(CH₃COO)₂ were purchased from E. Merck. All the chemicals were used without further purification. The ligand 4'-nitrobenzaldehyde-4-phenylsemicarbazone was prepared by refluxing the mixture of equimolar solution of nitrobenzaldehyde and 4-phenylsemicarbazide in ethanol in the presence of few drops of conc. H₂SO₄ for 1 h (**Scheme-**I). The solid obtained on cooling the solution was recrystallized in ethanol into yellowish white crystal with 85 % yield.

The other ligand *i.e.*, 4'-nitrobenzaldehyde-4-phenylthiosemicarbazone was prepared by the similar method of reflux like above wherein 4-phenylsemicarbazide was replaced by 4-phenylthiosemicarbazide. The crude solid was recrystallized



Scheme-I: Structure of 4'-nitrobenzaldehyde-4-phenyl semicarbazone (NBPS) and 4'-nitrobenzaldehyde-4-phenyl thiosemicarbazone (NBPTS)

in ethanol whereby light yellow crystal were obtained with 86 % yield.

Both ligands were used for complexation with Fe(II), Ni(II) and Cu(II) by the usual method of refluxing, the mixture of metal(II) acetate with ligands taken in 1:2 ratio in ethanol for 3-4 h. The precipitate was dried in desiccators on anhydrous calcium chloride. The yields were 74-80 %. Their melting point was recorded.

Physical and analytical measurements: C, H, N content of ligands and complexes was determined using Perkin-Elemer 2400 CHN Elemental Analyzer. Iron was estimated by volumetrically using N/10 K₂Cr₂O₇ solution. Copper was estimated iodometrically using standard sodium thiosulphate solution. Cobalt was estimated gravimetrically using α -nitroso β napthol. The molar conductivity of complexes was determined using CG-857 Schott-Grat Gambit Conductivity meter in DMF solution of 10⁻³ M molar concentration. The IR spectra of ligands as well as complexes were recorded on Perkin-Elemer FTIR-Spectrometer, Spectrum-two in range of 4000-400 cm⁻¹ using KBr Pellets. The electronic spectra of complexes were recorded on Perkin-Elmer UV visible Spectrometer Lambda-25 using DMSO solution. The magnetic susceptibility of complexes was determined on Gouy balance at room temperature using mercury tetrathiocynatocobaltate as calibrate.

RESULTS AND DISCUSSION

The percentage composition of ligands and complexes have been displayed in Table-1. On the basis of percentage composition and molar conductivity values, the complexes have been formulated as $[ML_2X_2]$ where M = Fe(II), Ni(II) and Cu(II), L = NBPS or NBPTS and X = CH₃COO⁻. The conductivity values of complexes are to low which indicate their non-electrolytic nature [16-19].

IR spectra of ligands and complexes: The IR spectra of free ligand 4'-nitrobenzaldehyde-4-phenyl semicarbazone a sharp band appears at 3150 cm⁻¹ which is assigned to v(N-H)[20]. It is further supported by a medium band at 1555 cm^{-1} due to bending vibration of δ (N–H). The strong band appearing at 3035 cm⁻¹ is due to v(C–H) stretching of benzene ring [21,22]. The medium band appearing at 2845 cm⁻¹ is due to v(C-H) of azomethine group. The strong band appearing at 1665 cm⁻¹ is fairly assigned to v(C=O) stretching of semicarbazone [23]. In the IR spectra of complexes of NBPS this band undergoes negative shift and appears at 1640 cm⁻¹ which is indicative of its coordination to the metal ions through oxygen [24,25]. The medium band appearing at 1630 cm⁻¹ in the IR spectra of free ligand is attributed to v(C=N) of azomethine group of the ligand [26]. In the IR spectra of complexes this band shifts to a lower frequency and appears at 1600 cm⁻¹. This shows the coordination of the ligand through azomethine nitrogen [27,28]. The medium and weak bands appearing at 1580, 1510 and 1455 cm⁻¹ are the characteristic absorption frequency of benzene ring vibration. The sharp band appearing at 1535 and 1340 cm⁻¹ may be assigned to $v_{asy}(NO_2)$ and $v_{sym}(NO_2)$ respectively. In the IR spectra of 4'-nitrobenzaldehyde-4-phenyl thiosemicarbazone the band near 1665 cm⁻¹ doesn't appear which shows the absence of C=O group in this ligand. This ligand absorbs strongly at 1230 cm⁻¹ due to v(C=S) stretching, which undergoes negative shift in its complexes. It is indicative of coordination through sulphur of C=S group of this ligand to the metal ions [29]. In the spectra of all the complexes two new bands appear at 1335 and 1330 cm⁻¹. These bands are the characteristic absorption bands for monodentataly coordinated CH₃COO⁻ ion to the metal ion in all the complexes. Thus on the basis of IR spectra of complexes in comparison to that of the free ligands, it may be asserted that the ligand NBPS acts as bidentate neutral ligand co-ordinating through carbonyl oxygen and azomethine nitrogen while the second ligand NBPTS also

TABLE-1									
COMPOSITION (%) FOUND (CALCD.)									
Compounds	Colour	Yield (%)	m.p. (°C)	М	С	Н	N	S	$\begin{array}{c} \lambda_{m}(\Omega^{-l}\\ cm^{2}mol^{-l}) \end{array}$
NBPS	Yellowish white	85	176	-	89.82	3.98	19.42	-	_
					(59.20)	(4.20)	(19.7)		
NBPTS	Light yellow	86	198	-	56.41	3.89	18.46	10.59	-
					(56.00)	(4.00)	(18.60)	(10.70)	
[Fe(NBPS) ₂ (CH ₃ COO) ₂]	Brilliant green	74	240	7.22	51.98	3.81	14.97	-	15
				(7.50)	(51.80)	(4.00)	(15.10)		
[Fe(NBPTS) ₂ (CH ₃ COO) ₂]	Dull green	80	242	6.98	49.91	3.72	14.29	8.12	16
				(7.20)	(49.60)	(3.90)	(14.50)	(8.30)	
[Ni(NBPS) ₂ (CH ₃ COO) ₂]	Dull red	80	240	7.90	51.76	3.91	14.87	-	13
				(8.00)	(51.50)	(4.00)	(15.00)		
[Ni(NBPTS) ₂ (CH ₃ COO) ₂]	Light red	78	246	7.43	49.68	3.72	14.28	8.12	15
				(7.60)	(49.40)	(3.90)	(14.40)	(8.20)	
[Cu(NBPS) ₂ (CH ₃ COO) ₂]	Bright blue	74	248	8.34	51.46	3.88	14.79	-	15
				(8.50)	(51.20)	(4.00)	(14.90)		
[Cu(NBPTS) ₂ (CH ₃ COO) ₂]	Blue	76	252	8.00	49.41	3.62	14.12	8.00	17
				(8.20)	(49.10)	(3.80)	(14.30)	(8.10)	

acts as bidentate neutral ligand coordinating through thiosulphur and azomethine nitrogen forming 5-membered chelate ring.

Magnetic moment and electronic spectra: The magnetic moment and electronic spectra of Fe(II) complexes have been given in Table-2.

TABLE-2							
Complexes	$v_{1} (cm^{-1})$	$v_{2} (cm^{-1})$	$v_{3} (cm^{-1})$	$\mu\left(BM\right)$			
[Fe(NBPS) ₂ (CH ₃ COO) ₂]	5600	9800	12800	5.08			
[Fe(NBPTS) ₂ (CH ₃ COO) ₂]	5400	9200	12600	5.07			

The magnetic moment of Fe(II) complexes are found to be 5.07 and 5.08 BM which correspond to high spin octahedral geometry to these complexes [30,31]. Fe(II) is a d^6 system for which the ground state term is ⁵D which splits into ⁵T_{2g} and ⁵E_g and thus only one band is expected in Fe(II) complexes. But here the complexes display three bands which indicates that the complexes have undergone tetragonal distortion (D_{4h}). This causes further splitting of T_{2g} in ⁵B_{2g} and ⁵E_g while, the excited state ⁵E_g splits into ⁵B_{1g} and ⁵A_{1g}. Thus these three bands may be assigned to ⁵B_{2g} \rightarrow ⁵E_g = v₁, ⁵B_{2g} \rightarrow ⁵B_{1g} = v₂ and ⁵B_{2g} \rightarrow ⁵A_{1g} = v₃ [32]. v₂ is the major of 10 Dq_{xy} while v₁ and v₃ involve also the other parameters like D_s and D_t. The values of different crystal field parameters are given in Table-3.

TABLE-3							
Complexes	$\begin{array}{c} \mathrm{Dq}_{\mathrm{xv}} \ (\mathrm{cm}^{-\mathrm{l}}) \end{array}$	$\begin{array}{c} \mathrm{Dq}_{z} \\ (\mathrm{cm}^{-1}) \end{array}$	D _s (cm ⁻¹)	$\begin{array}{c} D_{t} \ (cm^{-1}) \end{array}$			
[Fe(NBPS) ₂ (CH ₃ COO) ₂]	980	1650	1229	-383			
[Fe(NBPTS) ₂ (CH ₃ COO) ₂]	920	1490	1257	-326			

In these complexes the greater value of Dq_z than Dq_{xy} and negative value of D_t conspicuously indicate tetragonal compression along Z-axis in octahedral symmetry.

The magnetic moment and electronic spectra of Ni(II) complexes have been displayed in Table-4.

The magnetic moment of Ni(II) complexes are 3.35 and 3.40 BM at room temperature which are higher than μ_s for two unpaired electrons of Ni(II) octahedral complexes. This increase in magnetic moment may be attributed to strong spin orbit coupling under which $\mu_{eff} = \mu_{s.0.} (1-4\lambda/10Dq)$. As the spin orbit coupling constant λ for d^8 system is negative ($\lambda = -315$), $\mu_{eff.}$ becomes greater than $\mu_{s.0.}$ [33,34].

The electronic spectra of these complexes display four bands which may be assigned to ${}^{3}B_{1g} \rightarrow {}^{3}B_{2g} = \nu_{1}$, ${}^{3}B_{1g} \rightarrow {}^{3}E_{g}^{(a)} = \nu_{2}$, ${}^{3}B_{1g} \rightarrow {}^{3}A_{2g} = \nu_{3}$ and ${}^{3}B_{1g} \rightarrow {}^{3}E_{g}^{(b)} = \nu_{4}$. Here ν_{1} is the major of 10Dq and other transitions involve D_{s} and D_{t} parameters also. The values of various crystal field parameters are given in Table-5.

The appearance of four bands is indicative of appreciable tetragonal deviation from octahedral symmetry while greater value of Dq_z in respect of Dq_{xy} and negative value of D_t confirm the tetragonal compression along Z-axis [35-37].

TABLE-5							
Complexes	$\begin{array}{c} Dq_{xy} \\ (cm^{-l}) \end{array}$	$\begin{array}{c} \mathrm{Dq}_{\mathrm{Z}} \ (\mathrm{cm}^{-\mathrm{i}}) \end{array}$	$\begin{array}{c} D_{s} \ (cm^{-1}) \end{array}$	D_{r} (cm ⁻¹)			
[Ni(NBPS) ₂ (CH ₃ COO) ₂]	1155	1755	1290	-343			
[Ni(NBPTS) ₂ (CH ₃ COO) ₂]	1130	1782	1222	-373			

The magnetic moment and electronic spectral bands of Cu(II) complexes have been displayed in Table-6.

TABLE-6							
Complexes	$v_{1}(cm^{-1})$	$v_{2} (cm^{-1})$	(cm^{-1})	$\mu(BM)$			
[Cu(NBPS) ₂ (CH ₃ COO) ₂]	7200	18600	21000	1.96			
[Cu(NBPTS) ₂ (CH ₃ COO) ₂]	7000	18250	21100	1.98			

The magnetic moment of Cu(II) complexes are 1.96 and 1.98 BM at room temperature which indicate that the complexes are magnetically dilute with octahedral or distorted octahedral geometry [38,39].

The electronic spectra of Cu(II) complexes display three bands which may be assigned to ${}^{2}A_{1g} \rightarrow {}^{2}B_{1g} = v_{1}$, ${}^{2}A_{1g} \rightarrow {}^{2}B_{2g}$ = v_{2} and ${}^{2}A_{1g} \rightarrow {}^{2}E_{g} = v_{3}$. The various crystal field parameters derived from these spin allowed transition are given in Table-7.

	TABLE-7					
Complexes	$\begin{array}{c} \mathrm{Dq}_{\mathrm{xy}} \\ (\mathrm{cm}^{-\mathrm{l}}) \end{array}$	$\begin{array}{c} \mathrm{Dq}_{\mathrm{Z}} \\ (\mathrm{cm}^{-1}) \end{array}$	D _s (cm ⁻¹)	$\begin{array}{c} D_t \\ (cm^{-1}) \end{array}$		
[Cu(NBPS) ₂ (CH ₃ COO) ₂]	1140	2701	685	-892		
[Cu(NBPTS) ₂ (CH ₃ COO) ₂]	1125	27045	592	-926		

The appearance of three bands is clearly the consequence of tetragonal distortion in O_h symmetry while the greater value of Dq_z than Dq_{xy} with negative value of D_t show without doubt that there is compression along Z-axis of O_h symmetry [40,41].

Biological evaluation: The ligands as well as their metal complexes were screened against Gram-positive bacteria *Staphylococcus aureus* and Gram-negative bacteria *Escherichia coli* at concentration 50 and 100 ppm in DMSO and the results were compared with straptomycin as standard. The activity index of ligands as well as their metal complexes was derived by the expression:

Activity index = $\frac{\text{Inhibition zone by test compound (diameter)}}{\text{Inhibition zone by control (diameter)}} \times 100$

The results displayed in Table-8 clearly indicates that both ligands are biological active. It also indicates that the antibacterial activity gets enhanced after complexation. The activity increases with increasing concentration of ligands and complexes, however all the ligands and complexes show less antibacterial activity in comparison to the standard. The thio ligand and its metal complexes record greater activity than semicarbazone ligand and their complexes, which may be due to the presence of sulphur in place of oxygen in thiosemicarbazone. Copper complex with thiosemicarbazone shows

TABLE-4								
Complexes	$v_{1} (cm^{-1})$	$v_{2} (cm^{-1})$	$v_{3} (cm^{-1})$	$v_4 (cm^{-1})$	$\mu \left(BM\right)$			
[Ni(NBPS) ₂ (CH ₃ COO) ₂]	11550	14550	16000	24170	3.35			
[Ni(NBPTS) ₂ (CH ₃ COO) ₂]	11300	14560	16300	24100	3.40			

ANTIBACTERIAL SCREENING DATA OF LIGANDS AND COMPLEXES [DIAMETER OF INHIBITION ZONE (mm) AFTER 24 h INCUBATION]									
Compoundo		Staphylococc	us aureus			Escherich	ia coli		
Compounds	Diameter of i	nhibition zone	Activ	ity (%)	Diameter of i	inhibition zone	Activ	Activity (%)	
Concentration (ppm)	50	100	50	100	50	100	50	100	
NBPS	25	35	55.5	58	28	39	54	61.0	
NBPTS	27	38	60.0	63	30	42	58	66.0	
[Fe(NBPS) ₂ (CH ₃ COO) ₂]	28	36	62.0	60	34	45	65	70.0	
[Fe(NBPTS) ₂ (CH ₃ COO) ₂]	32	40	71.0	67	36	48	69	75.0	
[Ni(NBPS) ₂ (CH ₃ COO) ₂]	28	37	62.0	62	35	46	67	72.0	
[Ni(NBPTS) ₂ (CH ₃ COO) ₂]	34	41	75.5	68	37	50	71	78.0	
[Cu(NBPS) ₂ (CH ₃ COO) ₂]	38	45	84.0	75	40	49	77	76.5	
[Cu(NBPTS) ₂ (CH ₃ COO) ₂]	40	49	89.0	82	46	54	88	84.0	
Streptomycin (standard)	45	60	100.0	100	52	64	100	100.0	

TABLE-8

the greatest activity than all other complexes which may be due to a toxic nature of copper metal. The increases in antibacterial activity of ligands after complexation may be attributed to the formation of chelates in complexes [12,22].

Conclusion

The study reveals that the ligands are weakly coordinated to the metal ions in XY plane than acetate ions along Z-axis due to the presence of strongly electron withdrawing-NO₂ group in planar ligands. This makes tetragonally distorted octahedral complexes with appreciable compression along Zaxis which is well supported by the higher values of $Dq_{(z)}$ than $Dq_{(xy)}$ and negative value of D_t in all the complexes. The antibacterial screening of complexes reveals that all the complexes have greater antibacterial activity than the free ligands confirming the concept of enhancement of antibacterial activities of a compound after complexatation.

REFERENCES

- 1. I.B. Bersuker, Jahn-Teller Effect and Vibronic Interaction in Modern Chemistry, Plenum Press, New York (1984).
- 2 D. Reinen and M. Atansov, Mag. Reson. Rev., 5, 167 (1971).
- 3. B.N. Figgis, Introduction to Ligand Field, Wiley Eastern Limited, New Delhi, pp. 212-214 (1964).
- W. Zhang, J.L. Loebach, S.R. Wilson and E.N. Jacobsen, J. Am. Chem. 4. Soc., 112, 2801 (1990);
- https://doi.org/10.1021/ja00163a052 F. Yamakura, K. Kobayashi, H. Ue and M. Konno, Eur. J. Biochem., 5. **227**, 700 (1995):
- https://doi.org/10.1111/j.1432-1033.1995.tb20191.x.
- 6. A. Bencini, I. Ciofini and M.G. Uytterhoeven, Inorg. Chim. Acta, 274, 90 (1998);
- https://doi.org/10.1016/S0020-1693(97)05985-9.
- J.P. Renault, C. Verchère-Béaur, I. Morgenstern-Badarau, F. Yamakura 7. and M. Gerloch, Inorg. Chem., 39, 2666 (2000); https://doi.org/10.1021/ic0000451.
- 8. S. Mossin, H. Weihe and A.-L. Barra, J. Am. Chem. Soc., 124, 8764 (2002); https://doi.org/10.1021/ja012574p.
- S.N. Pandeya, P. Yogeeswari and J.P. Stables, Eur. J. Med. Chem., 35, 9. 879 (2000);

https://doi.org/10.1016/S0223-5234(00)01169-7.

- 10 M.R.P. Kurup, B. Varghese, M. Sithambaresan, S. Krishnan, S.R. Sheeja and E. Suresh, Polyhedron, 30, 70 (2011); https://doi.org/10.1016/j.poly.2010.09.030.
- 11. V.L. Siji, M.R. Sudarsanakumar and S. Suma, Transition Met. Chem., 36, 417 (2011);

https://doi.org/10.1007/s11243-011-9485-z.

12. S.R. Layana, V.L. Sigi, M.R. Sudarasana Kumar, S. Kumar, M.R.P. Kurup and T.S. Sikha, J. Indian Chem. Soc., 93, 577 (2016).

- 13. V.S. Shekhawat, S. Varshney and A.K. Varshney, J. Indian Chem. Soc., 94, 21 (2017).
- 14. S.A. Khan and N. Yusuf, Eur. J. Med. Chem., 44, 2270 (2009); https://doi.org/10.1016/j.ejmech.2008.06.008
- 15. A.E. Graminha, C. Rodrigues, A.A. Batista, L.R. Teixeira, E.S. Fagundes and H. Beraldo, Spectrochim. Acta A Mol. Biomol. Spectrosc., 69, 1073 (2008);
 - https://doi.org/10.1016/j.saa.2007.06.005.
- B.B. Mahapatra and S.K. Saraf, J. Indian Chem. Soc., 80, 696 (2003). 16.
- 17. S.K. Sinha, D. Kumar, C.M. Chandra, B.N. Subash, S. Kumar and S. Sharma, Int. J. Develop. Res., 7, 12295 (2017).
- 18. A. Sheela and N.L.H. Nair, J. Indian Chem. Soc., 89, 445 (2012).
- R. Ranjan, N. Sinha, S. Kumar, C. M. Chandra and S. Sharma, IRA-19. Int. J. Appl. Sci., 7, 34 (2017).
- D. Das, A.K. Panda, P. Jana, S.B. Patjoshi and A. Mahapatra, J. Indian Chem. Soc., 79, 48 (2002).
- 21 M.K. Singh, A.D.R. Laskar and B. Pant, J. Indian Chem. Soc., 85, 485 (2008).
- 22. A. Karim, R. Rani, R. Ranjan, U. Kumar, V. Kumar and S. Sharma, Asian J. Chem., 29, 626 (2017);
- https://doi.org/10.14233/ajchem.2017.20286. S. Singh, P. Tripathi, O.P. Pandey and S.K. Sen Gupta, J. Indian Chem. 23.
- Soc., 90, 565 (2013).
- B.K. Rai and R. Kumari, Asian J. Chem., 23, 4625 (2011). 24.
- 25. N.P. Singh, K.L. Anand and J. Singh, Asian J. Chem., 23, 4090 (2011).
- 26. R. Ranjan, R. Rani, S.S. Singh, A.K. Singh Jr. and S. Charma, Asian J. Chem., 22, 7580 (2010).
- A.K. Singh, S.P. Sinha, V. Kumar, S.S. Rai and S. Sharma, Asian J. 27. Chem., 23, 43 (2011).
- 28. R.L. De, R.K. De, I. Banerjee, N. Mandal, N. Ray, J. Mukharjee and E. Keller, J. Indian Chem. Soc., 90, 27 (2013).
- G.S. Sanya, P.K. Nath and R. Ganguly, J. Indian Chem. Soc., 79, 54 (2002). 29 M.A. Halcrow, Coord. Chem. Rev., 253, 2493 (2009); 30.
- https://doi.org/10.1016/j.ccr.2009.07.009.
- M. Matsuda, H. Isozaki and H. Tajima, Chem. Lett., 37, 374 (2008); 31. https://doi.org/10.1246/c1.2008.374.
- 32. A.B.P. Lever, Inorganic Electronic Spectroscopy, Elsevier, New York, edn 2, p. 19 (1984).
- 33 P. Bhatnagar and N.K. Bhatnagar, J. Indian Chem. Soc., 89, 1273 (2012).
- 34. N. Agrawal and A. Singh, J. Indian Chem. Soc., 90, 585 (2013).
- 35. M.A. Hitchman, Inorg. Chem., 11, 2387 (1972); https://doi.org/10.1021/ic50116a019. 36. E.I. Solomon and C.J. Ballhausen, Mol. Phys., 29, 279 (1975);
- https://doi.org/10.1080/00268977500100191
- 37. A.B.P. Lever, G. London and P.J. McCarthy, Can. J. Chem., 55, 3172 (1977):

https://doi.org/10.1139/v77-445.

- B.B. Mahapatra and N. Patel, J. Indian Chem. Soc., 86, 518 (2009). 38
- D. Kumar, S. Sharma and R.C. Sharma, J. Indian Chem. Soc., 87, 1547 39. (2010).
- M. Fujiwara, Y. Nakajima, T. Matsushita and T. Shumo, Polyhedron, 40. 4, 1589 (1985);

https://doi.org/10.1016/S0277-5387(00)87233-2.

41. R.N. Patel, A.P. Patel and K.B. Pandaya, J. Indian Chem. Soc., 78, 6, (2001).